

# Differential Geometry 1999. Exercises.

Michael Murray

September 18, 2002

## Contents

1	Co-ordinate charts and manifolds.	1
2	Smooth functions.	2
3	Submanifolds.	3
4	Vector fields and differential forms.	3
5	Complex line bundles	4

## 1 Co-ordinate charts and manifolds.

*Exercise 1.1.* Consider the  $n$  sphere

$$S^n = \{x \in \mathbb{R}^{n+1} \mid \|x\|^2 = 1\}.$$

Let

$$U_i = \{x \in S^n \mid x^i \neq 1\} = S^n - \{e_i\}$$

where  $e_i$  has all components 0 except the  $i$ th which is equal to one. If  $x$  is a point in  $U_i$  show that there is a unique line through  $x$  and the vector  $e_i$ . Show that this line intersects the plane

$$\{x \mid x^i = 0\}$$

in exactly one point. Writing this point as

$$(\psi_i^1(x), \psi_i^2(x), \dots, \psi_i^{i-1}(x), 0, \psi_i^i(x), \dots, \psi_i^n(x))$$

defines a function

$$\psi_i = (\psi_i^1, \dots, \psi_i^n): U_i \rightarrow \mathbb{R}^n.$$

Show that  $(U_i, \psi_i)$  is a co-ordinate chart on  $S^n$  and that

$$\{(U_i, \psi_i) \mid i = 1, \dots, n\}.$$

is an atlas for  $S^n$ .

The functions  $\psi_i$  are said to arise by *stereographic projection* from  $e_i$  onto the plane  $\{x \mid x^i = 0\}$ .

*Exercise 1.2.* Consider the sphere  $S^n$  again. Define

$$U_i^+ = \{x \in S^n \mid x^i > 0\}$$

and define  $\psi_i^+ : U_i^+ \rightarrow \mathbb{R}^n$  by

$$\psi_i^+(x^1, \dots, x^{n+1}) = (x^1, \dots, x^{i-1}, x^{i+1}, \dots, x^n).$$

Show that  $(U_i^+, \psi_i^+)$  is a co-ordinate chart for  $S^n$ . Similarly define

$$U_i^- = \{x \in S^n \mid x^i < 0\}$$

and define  $\psi_i^- : U_i^- \rightarrow \mathbb{R}^n$  by

$$\psi_i^-(x^1, \dots, x^{n+1}) = (x^1, \dots, x^{i-1}, x^{i+1}, \dots, x^n).$$

Again show that  $(U_i^-, \psi_i^-)$  is a co-ordinate chart for  $S^n$ .

Show that

$$\{(U_i^+, \psi_i^+), (U_i^-, \psi_i^-) \mid i = 1, \dots, n\}$$

is an atlas for  $S^n$ .

*Exercise 1.3.* Show that the atlases in Exercises 1.1 and 1.2 define the same maximal atlas on  $S^n$ .

*Exercise 1.4.* Let  $\mathbb{R}P_n$  be the set of all lines (through the origin) in  $\mathbb{R}^{n+1}$ . This space is called real, projective space of dimension  $n$ . If  $x$  is a non-zero vector in  $\mathbb{R}^{n+1}$  denote by  $[x]$  the line through  $x$ . Show that  $[x] = [y]$  if and only if there is a non-zero real number  $\lambda$  such that  $x = \lambda y$ .

Define subsets  $U_i$  of  $\mathbb{R}P_n$  by

$$U_i = \{[x^0, \dots, x^n] \mid x^i \neq 0\}$$

and maps  $\varphi_i : U_i \rightarrow \mathbb{R}^n$  by

$$\varphi_i([x^0, \dots, x^n]) = \left( \frac{x^0}{x^i}, \dots, \frac{x^{i-1}}{x^i}, \frac{x^{i+1}}{x^i}, \dots, \frac{x^n}{x^i} \right)$$

for every  $i = 0, \dots, n$ . Show that  $\varphi_i$  is well defined and that  $(U_i, \varphi_i)$  is a co-ordinate chart on  $\mathbb{R}P_n$ . Show that

$$\{(U_i, \varphi_i) \mid i = 0, \dots, n\}$$

is an atlas for  $\mathbb{R}P_n$ .

*Exercise 1.5.* Show that if  $M_1$  and  $M_2$  are manifolds then there is a natural way of making  $M_1 \times M_2$  into a manifold so that  $\dim(M_1 \times M_2) = \dim(M_1) + \dim(M_2)$ .

*Exercise 1.6.* Repeat exercise (1.4) for  $\mathbb{C}^n$  to define  $2n$  dimensional complex projective space  $\mathbb{C}P_n$  as the space of complex lines through zero in  $\mathbb{C}^{n+1}$ .

## 2 Smooth functions.

*Exercise 2.1.* Define  $h : \mathbb{R} \rightarrow \mathbb{R}$  by

$$h(x) = \begin{cases} \exp\left(\frac{-1}{1-x^2}\right) & \text{if } -1 < x < 1, \\ 0 & \text{otherwise.} \end{cases}$$

and show that  $h$  is smooth. By integrating  $h$  find a smooth function  $g : \mathbb{R} \rightarrow \mathbb{R}$  with the property that  $g(x)$  is zero for  $x < -1$  and  $g(x)$  is one for  $x > 1$ . Show that for any  $\epsilon > \delta > 0$  there is a smooth function  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$  with  $\phi(x)$  equal to zero if  $\|x\| > \epsilon$  and  $\phi$  equal to one if  $\|x\| < \delta$ . Now consider a manifold  $M$  and a point  $x$ . By using co-ordinates show that if  $U$  is any open subset of  $M$  containing  $x$  then there are open subsets  $U_1$  and  $U_2$  with  $x \in U_1 \subset U_2 \subset U$  and a smooth function  $f : M \rightarrow \mathbb{R}$  with  $f$  equal to 1 on all of  $U_1$  and equal to zero outside of  $U_2$ .

*Exercise 2.2.* Let  $x$  be point in a manifold  $M$ . Let  $X_x$  be the set of all pairs  $(U, f)$  where  $U$  is a open set containing  $x$  and  $f : U \rightarrow \mathbb{R}$  is a smooth function. Define a relation on  $X_x$  by saying that  $(U, f) \simeq (V, g)$  if there is an open set  $W$  with  $x \in W \subset U \cap V$  and  $f|_W = g|_W$ . Show that this an equivalence relation. Equivalence classes are called *germs* at  $x$  and the set of them we will denote by  $G_x$ . Show that  $G_x$  is an algebra under pointwise addition, scalar multiplication and multiplication. If  $f \in C^\infty(M, \mathbb{R})$  the algebra of all smooth functions on  $M$  it defines the germ containing  $(M, f)$ . Show that the map this induces  $C^\infty(M, \mathbb{R}) \rightarrow G_x$  is onto. [Hint: Use 2.1.]

*Exercise 2.3.* Consider the map  $F: \mathbb{R}^3 \rightarrow \mathbb{R}^2$  defined by

$$F(x, y, z) = (x^2 + y^2 + z^2 - 9, x + y + z - 3).$$

If we identify the tangent spaces to  $\mathbb{R}^3$  and  $\mathbb{R}^2$  with  $\mathbb{R}^3$  and  $\mathbb{R}^2$  respectively calculate the tangent map

$$T_{(x,y,z)}F: \mathbb{R}^3 \rightarrow \mathbb{R}^2.$$

*Exercise 2.4.* Define a map  $F: S^2 \rightarrow \mathbb{C}P_1$  by

$$F(x, y, z) = [x + iy, 1 - z].$$

By using the co-ordinates defined in Exercises (1.1) and (1.6) show that this map is well defined as  $z \rightarrow 1$  and that it is, in fact, a diffeomorphism.

### 3 Submanifolds.

*Exercise 3.1.* Show that the set defined by the equation

$$r^2 - a^2 = (\sqrt{x^2 + a^2} - a)^{1/2}$$

is a smooth submanifold of  $\mathbb{R}^3$  if  $a$  and  $r$  are real numbers with  $r < a$ .

*Exercise 3.2.* Show that the following subset of  $\mathbb{R}^3$  is a submanifold:

$$Q = \{(x, y, z) \mid x^2 + y^2 + z^2 = 9 \text{ and } x + y + z = 3\}.$$

### 4 Vector fields and differential forms.

*Exercise 4.1.* Let  $X$  and  $Y$  be vector fields on a manifold  $M$ . Define a new vector field  $[X, Y]$  by defining it in local co-ordinates  $(U, \phi)$  by

$$[X, Y]_{|U} = \sum_{i,j} (X_i \frac{\partial^j}{\partial \phi^i} \partial \phi^j - Y_j \frac{\partial X^j}{\partial \phi^i}) \frac{\partial}{\partial \phi^j}.$$

Show that this makes sense. That is it doesn't really depend on the choice of co-ordinates. The vector field  $[X, Y]$  is called the Lie bracket of  $X$  and  $Y$ .

*Exercise 4.2.* If  $X$  is a vector field and  $\omega$  is a differential 1-form show that the differential 1-form defined by

$$L_X(\omega) = \sum_{i,j} (X^i \frac{\partial \omega_j}{\partial \theta^i} + \omega_i \frac{\partial X^i}{\partial \theta^j}) d\theta^j.$$

where

$$X = \sum_i X^i \frac{\partial}{\partial \theta^i} \quad \text{and} \quad \omega = \sum_i \omega_i d\theta^i$$

is actually independent of the choices of co-ordinates. We call  $L_X(\omega)$  the Lie derivative of  $\omega$  by  $X$ .

*Exercise 4.3.* Let  $\alpha$  and  $\beta$  be  $p$  and  $q$  forms, respectively on a manifold  $M$ . Show that

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^p \alpha \wedge d\beta.$$

*Exercise 4.4.* Consider the circle  $S^1 = \{(x, y) \mid x^2 + y^2 = 1\}$ . This is a manifold of dimension 1. The circle has a co-ordinate chart  $(U, \theta)$  where  $U = S^1 - \{(1, 0)\}$  and  $\theta: U \rightarrow (0, 2\pi)$  is defined implicitly by

$$(x, y) = (\cos(\theta(x, y)), \sin(\theta(x, y))).$$

That is  $\theta$  is the usual angle co-ordinate in polar co-ordinates. Identify the tangent space to the circle at  $(x, y)$  with the line in  $\mathbb{R}^2$  tangential to the circle at  $(x, y)$ . Calculate a formula for the vector field  $\partial/\partial\theta$  in terms of  $x$  and  $y$  and hence show that it extends from  $U$  to a vector field on all of  $S^1$ . Show that  $d\theta$  also extends to a differential 1-form  $\omega$  on all of the circle. Show that there is no function  $f: S^1 \rightarrow \mathbb{R}$  such that  $\omega = df$ .

*Exercise 4.5.* Let  $S^2 = \{x \in \mathbb{R}^3 \mid \|x\|^2 = 1\}$  be the two-sphere. Recall that the spherical co-ordinates  $(\theta, \phi)$  of the point  $(x, y, z)$  on the two-sphere are defined by requiring that:

$$\begin{aligned}x &= \sin(\psi) \cos(\theta) \\y &= \sin(\psi) \sin(\theta) \\z &= \cos(\psi).\end{aligned}$$

Find an open set  $U \subset S^2$  for the domain of the spherical co-ordinates so that  $\psi \in (0, \pi)$  and  $\theta \in (0, 2\pi)$ .

For any  $x$  in  $S^2$  and  $X, Y \in T_x S^2$  define a differential two-form  $\omega$  on  $S^2$  by  $\omega_x(X, Y) = \langle x, X \times Y \rangle$ , where  $\langle \cdot, \cdot \rangle$  is the usual inner-product on  $\mathbb{R}^3$  and  $x$  is the cross-product of three vectors. By using suitable co-ordinates (spherical are good) calculate the integral of  $\omega$  over  $S^2$  and show that it is non-zero.

*Exercise 4.6.* Show that it is not possible to find a differential one-form  $\mu$  on the two sphere such that  $d\mu$  is the volume form  $\omega$  defined in exercise (4.5).

*Exercise 4.7.* Consider the torus  $T^2$  in  $\mathbb{R}^3$  with co-ordinates  $(\theta, \phi)$  defined implicitly by

$$x = (b + a \sin(\phi)) \cos(\theta), (b + a \sin(\phi)) \sin(\theta), a \cos(\phi).$$

Calculate  $\partial/\partial\psi$  and  $\partial/\partial\theta$ . Calculate the (outward) unit normal  $n(x)$  to the torus, this is the vector in  $\mathbb{R}^3$  orthogonal to the tangent space to the torus at  $x$ . You will need to draw a picture or something to check it is the outward normal.

Define  $\text{vol}$  a two-form by  $\text{vol}(X, Y) = \langle n, X \times Y \rangle$  and calculate its integral over  $T^2$  when we orient  $T^2$  in such a way as to make  $\text{vol}$  positive.

*Exercise 4.8.* Recall the definition of  $\mathbb{R}P_2$  the space all lines through the origin in  $\mathbb{R}^2$  and its associated co-ordinate charts given in Exercise 1.4. Calculate the linear relationship between the basis of one forms  $d\psi_i^1, d\psi_i^2$  and the basis of one forms  $d\psi_j^1, d\psi_j^2$  for  $i \neq j$ . Hence calculate the relationship between  $d\psi_i^1 \wedge d\psi_i^2$  and  $d\psi_j^1 \wedge d\psi_j^2$ . Show that  $\mathbb{R}P_2$  is not orientable.

*Exercise 4.9.* Let  $f: M \rightarrow N$  be a smooth map. If  $\omega$  is a  $p$ -form on  $N$  show that  $df^*(\omega) = f^*d\omega$ .

## 5 Complex line bundles

*Exercise 5.1.* Let  $\nabla^0$  and  $\nabla^1$  be connections on a complex line bundle  $L$  and define

$$\nabla^t(\phi) = t\nabla^1(\phi) + (1-t)\nabla^0(\phi)$$

for any section  $\phi$  of  $L$ . Show that  $\nabla^t$  is a connection for any real number  $t$ . Calculate its curvature.

*Exercise 5.2.* Show that if  $L \rightarrow M$  is a trivial bundle then it has zero Chern class.

*Exercise 5.3.* Consider the Hopf bundle  $H$  over  $\mathbb{C}P_1$ . Define parameters on  $U_0 = \mathbb{C}P_1 - [1, 0]$  by  $(x, y) \mapsto [x + iy, 1]$ . Let  $s_0([x + iy, 1]) = ([x + iy, 1], (x + iy, 1))$  be the section defined in class. Using (hermitian) orthogonal projection define a connection  $\nabla$  on  $H$  and calculate the connection one form  $A_0$ . Be careful to make the orthogonal projection complex linear. Calculate the curvature over the open set  $U_0$  and integrate it over  $U_0$  to find the Chern class of  $H$ . You may find it convenient to work with the complex differential forms  $dz = dx + idy$  and  $d\bar{z} = dx - idy$ .

*Exercise 5.4.* Consider the tangent bundle to the two-sphere. Give it the connection defined by orthogonal projection and calculate its curvature and hence the chern class of the tangent bundle to the two-sphere.

*Exercise 5.5.* Repeat Exercise 5.4 for the torus using the co-ordinates defined in Exercise 4.7.

*Exercise 5.6.* This assumes you are familiar with the Gauss-Bonnet theorem. If  $\Sigma$  is a closed surface in  $\mathbb{R}^3$  define a connection on its tangent bundle by using orthogonal projection. Relate the curvature of this connection to the usual Gaussian curvature.